

Original Article

Gastroprotective effect of sodium hydrosulfide against indomethacin-induced gastric ulcer in diabetic rats

Nasim Nazariani¹, Seyyed Ali Mard^{2*} , Sima Nasri¹, Ali Veisi³

1. Biology Department, Payame Noor University, Tehran, Iran

2. Alimentary Research Center, Physiology Research Center (PRC), Department of Physiology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

3. Behbahan Faculty of Medical Sciences, Behbahan, Iran

Abstract

Introduction: The incidence rate of gastric erosions and ulcers in diabetic patients are higher due to failure of mucosal antioxidant defense and maintain enough blood flow. The present study evaluated the gastro-protective effect of sodium hydrosulfide (NaHS) against indomethacin-induced gastric lesions in diabetic rats.

Methods: In order to test anti-ulcer activity of NaHS against indomethacin, four diabetic groups of rats including diabetic control and 3 NaHS-treated groups received a single dose of physiologic saline or NaHS at 320, 640 and 1280 µg/kg respectively, 30 min before ulcer induction by indomethacin. Five hours later, the animals were killed and their stomachs were removed for macroscopically and microscopically evaluations. In order to evaluate the antacid effect of NaHS, 4 groups of diabetic rats received physiologic saline or NaHS at 320, 640 and 1280 µg/kg and 30 min later anesthetized, underwent a midline laparotomy and then their pylorus ligated. Five hours later, the animals were killed, their stomachs were removed and pH of gastric effluents were measured.

Results: Indomethacin induced gastric lesions in glandular part of the stomach. NaHS at 640 and 1280 µg/kg significantly decreased the indomethacin-induced gastric lesions in diabetic rats. The pH of gastric effluents and mucus content increased by NaHS at doses of 640 and 1280 µg/kg. Macroscopic and microscopic observations showed that mucosal erosions induced by indomethacin were significantly inhibited by NaHS.

Conclusion: results suggest NaHS through decreasing the rate of gastric acid output and increasing the mucus production, protected the gastric mucosa against indomethacin-induced gastric lesions in diabetic rats.

Keywords:

Diabetes;
Gastric acid secretion;
Sodium hydrosulfide;
Rat

Received: 23 Aug 2017

Accepted: 3 Jan 2018

*Correspondence to:

S.A. Mard

Tel: +98-6133362411

Email:

mard-sa@ajums.ac.ir

Introduction

The gastro-protective activity of sodium hydrosulfide (NaHS) well established against different gastric irritants such as ischemia-reperfusion injury, non-

steroidal anti-inflammatory drugs (Wallace, 2012) and water-immersion restraint stress (Magierowski et al., 2015). Studies have been shown that NaHS through a variety of mechanisms including inhibition of the acid output in response to distention and histamine, increasing the mucosal nitric oxide level (Mard et al.,

2014), promoting the production of mucous secretion and increasing the mRNA expression level of prostaglandin E2 receptors (Mard et al., 2017) contributed in mucosal defense barrier.

It has been shown that the gastric mucosa in diabetic rats has a more susceptibility to gastric irritants (Takeuchi et al., 1994; Tashima et al., 1998). A study showed that the prevalence of gastric erosions and gastric ulcers in diabetic patients is more than 45% (Boehme et al., 2007). Moreover, according to twenty-four pH studies, the prevalence of asymptomatic reflux in insulin-dependent diabetic patients were 28% higher than in healthy population (Lluch et al., 1999). There are many reasons for decreased ability of gastric mucosa against irritants such as inability of the gastric mucosa to deliver the enough amount of calcitonin-gene related peptide, impairment of gastric mucosal blood flow and decrement the efficiency of antioxidant defense system in the gastric mucosal layer (Goldin et al., 1997; Tashima et al., 1998). Therefore, the present study was designed to evaluate the gastro-protective effect of NaHS against indomethacin-induced gastric lesions in diabetic rats.

Materials and methods

Animals

Male Wistar rats weighing 150-200 g were supplied from the animal house of Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. Animals were fed on conventional diets and tap water. They were maintained under standard conditions of humidity, temperature (22±2 °C) and light/dark cycle (12:12 h). Animals were fasted 24 h before the experiments.

Animal grouping and procedures

In the first set of experiments, to evaluate the protective effect of NaHS on indomethacin-induced gastric lesions, four diabetic groups of rats were used. They were diabetic control group and 3 NaHS-treated groups (6 in each group). The control rats were given an intraperitoneally (ip) injection of normal saline (2 ml/kg) and NaHS-treated groups received a single ip injection of NaHS (Sigma, USA) at doses 320, 640 or 1280 µg/kg. Thirty min after administration of normal saline or NaHS, all groups of animals received indomethacin (40 mg/kg, orally) to

induce gastric lesions (Kianbakht and Mozaffari, 2009). Five hour after the administration of indomethacin, animals sacrificed under a high dose of anesthetics (100 mg/kg ketamine + 20 mg/kg xylazine, ip, Alfasan Co.Woerden-Holland), their stomachs rapidly removed, opened along the greater curvature, rinsed with physiologic saline and pinned out on ice-cooled saline.

Evaluation of effects of NaHS on gastric mucus production

To evaluate the mucus production, gastric mucus was gently scraped using a glass slide and the obtained mucus weighed with an accurate digital device.

Evaluation of NaHS effects on gastric acid secretion

To determine the antisecretory effect of NaHS, 4 separate groups of fasted rats (n=5 in each group) were subjected to a pylorus ligation procedure as described by Shay et al. (1945). The control and 3 NaHS-treated groups received a single ip injection of saline or NaHS at doses of 320, 640 or 1280 µg/kg, respectively, 30 min prior to pylorus ligation. Five hours after pylorus ligation, animals were killed by an overdose of anesthetics (ketamine+xylazine), their stomachs removed and the entire gastric contents were transferred into centrifuge tubes for measuring pH using a digital pH meter (iSTEK, South Korea).

Induction of diabetes

The fasted rats were received a single ip injection of alloxan monohydrate (Sigma, USA) at 130 mg/kg. After 72 h, the level of fasting blood glucose was checked using a glucometer (Elegance, Frankenberg, Germany). Rat with fasting blood glucose higher than 250 mg/dl was considered as diabetic. Two weeks later, animals with fasting blood sugar more than 250 mg/dl subjected to induce the gastric ulcer (Mard et al., 2016; Qiu et al., 2008).

Histopathological evaluation

For histological evaluation, stomachs from control and treated animals were fixed in 10% formalin, dehydrated in grade ethanol and embedded in paraffin. Thereafter, sections of tissue were cut at 5 µm using a microtome, stained with hematoxylin and eosin and assessed under an Olympus microscope




Fig.1. Histological evaluation of the gastric mucosa. Microscopic representation of the gastric sections 5 hour after indomethacin administration in control group (A) shows multiple erosions, exfoliation and necrosis of superficial cells as well as hemorrhage in the mucosal layer of the gastric mucosa. Severe hemorrhagic and necrotic areas in Fig.1B shows that NaHS at dose of 320 $\mu\text{g}/\text{kg}$ was not enough to protect the gastric mucosa against indomethacin; however NaHS at doses 640 and 1280 $\mu\text{g}/\text{kg}$ (C and D) improved the histopathological changes. All sections stained with hematoxylin and eosin; $\times 100$ magnification.

(IX50).

Statistical analysis

Data are shown as mean \pm SEM. Statistical analysis was performed by one-way ANOVA and followed by post hoc Tukey's Test. Significance was set at a $P < 0.05$ level.

Results

Effect of NaHS on indomethacin-induced morphological and histological alteration of the gastric mucosa in diabetic rats

As shown in Figure 1, oral administration of indomethacin caused multiple mucosal erosions, exfoliation and necrosis of superficial cells, and hemorrhages in the mucosal layer in the gastric mucosa (Fig. 1A). The microscopic and macroscopic

data showed that the lowest studied dose of NaHS (320 $\mu\text{g}/\text{kg}$) could not prevent the gastric mucosa against indomethacin-induced gastric lesions (Fig. 1B). Pretreatment with NaHS at doses 640 and 1280 $\mu\text{g}/\text{kg}$ effectively improved the histopathological changes (Figs. 1C and D). Figure 2 shows that the total area of gastric lesions in NaHS-treated rats were lower than in control group. The area of gastric lesions in groups received NaHS at doses 640 and 1280 $\mu\text{g}/\text{kg}$ was significantly lower than in control rats ($P < 0.05$ and $P < 0.01$, respectively)

Effect of NaHS pretreatment on the mucus production of gastric wall in indomethacin-induced gastric ulcer rats

As shown in Figure 3, pretreatment with NaHS increased the rate of mucus production in indomethacin-induced gastric lesion rats. The weight




Fig.2. The effect of sodium hydrosulfide on indomethacin-induced gastric ulcer in diabetic rats. NaHS at doses 640 and 1280 $\mu\text{g}/\text{kg}$ decreased the total area of gastric ulcer. C: diabetic control rats; D+NaHS-(320), (640) or (1280): diabetic rats pretreated with NaHS at 320, 640 or 1280 $\mu\text{g}/\text{kg}$. * $P<0.05$ and ** $P<0.01$ versus diabetic control group.




Fig. 3. The effect of sodium hydrosulfide on the production of gastric wall mucus. NaHS at doses of 640 and 1280 $\mu\text{g}/\text{kg}$ increased the rate of mucus secretion by the gastric mucosa. C: diabetic control rats; D+NaHS-(320), (640) or (1280): diabetic rats pretreated with NaHS at 320; 640 or 1280 $\mu\text{g}/\text{kg}$. * $P<0.05$ and *** $P<0.001$ versus diabetic control group.




Fig. 4. The effect of sodium hydrosulfide on the gastric acid output in pylorus-ligated rats. NaHS at doses 320 and 1280 $\mu\text{g}/\text{kg}$ decreased the total acid content in the gastric effluents. C-PL: control pylorus-ligated rats; PL: pylorus ligated rats. The (320), (640) and (1280) shows the given dose of NaHS in each group. * represents $P < 0.05$ compared to control.

of mucus in NaHS-treated diabetic rats received NaHS at doses 640 or 1280 $\mu\text{g}/\text{kg}$ was significantly higher than in control rats ($P < 0.001$ and $P < 0.05$, respectively).

Effect of NaHS on the profile of gastric secretion induced by pylorus-ligated rats

As shown in Figure 4, the total acid content of the gastric effluents significantly decreased by NaHS at doses 320 and 1280 $\mu\text{g}/\text{kg}$ ($P < 0.05$ at both cases).

Discussion

The results of the present study showed that a single administration of NaHS in diabetic rats: 1. decreased the total area of gastric lesions induced by indomethacin; 2. decreased the gastric acid output and 3. increased the rate of mucus production in the gastric wall.

Our results showed that NaHS at 640 and 1280 $\mu\text{g}/\text{kg}$ had a stimulatory effect on the production of gastric wall mucus in diabetic rats. It is a general fact that prostaglandins E_2 and $F_{2\alpha}$ increase the rate of mucous production/release in gastric mucosal layer. A study showed that exogenous H_2S by increasing the production of prostaglandin E_2 protected the gastric mucosa against water-immersion restraint stress in rats (Magierowski et al., 2015). Another recently published research showed that NaHS up-regulated the mRNA expressions of prostaglandin E_2

receptors in male Wistar rats in response to mucosal acidification by 100 mM hydrochloric acid and distention-induced gastric acid secretion (Mard et al., 2017). These finding together showed that NaHS by increasing the levels of prostaglandin E_2 and/or up-regulating its receptors, increased the production of gastric wall mucous in diabetic rats received NaHS. Moreover, these finding was in agreement with the previous report as shown by authors NaHS at dose 80 $\mu\text{g}/\text{kg}$ increased this rate in response to distention-induced acid output in nondiabetic rats (Mard et al., 2017). As shown in the current results, NaHS at 320 $\mu\text{g}/\text{kg}$ failed to stimulate the mucus secretion in diabetic rats. This finding implied that the needed dose of NaHS to increase the rate of mucus secretion in diabetic rats is more than in the nondiabetic ones. However, the results showed that this effect was not dose-dependent because the highest response was not observed in rats received the highest dose of NaHS. As shown in the present results, NaHS at 640 $\mu\text{g}/\text{kg}$ had the strongest stimulatory effect on the mucus secretion. This finding seems to be a secondary/compensatory response to lower potency of this dose to inhibit the acid output compared to the rats received NaHS at 1280 $\mu\text{g}/\text{kg}$. Moreover, the current results showed that NaHS at 1280 $\mu\text{g}/\text{kg}$ effectively decreased the acid output while this effect was not observed in rats received it at 640 $\mu\text{g}/\text{kg}$. Therefore, comparing antiulcer effect of these two effective doses (640 or 1280 $\mu\text{g}/\text{kg}$) showed that

inhibiting the acid output plus stimulating the production of gastric wall mucus as occurred in response to NaHS at 1280 µg/kg had a stronger antiulcer effect.

The antacid result of NaHS at 320 µg/kg was not enough to inhibit/decrease the inducing of gastric lesions by indomethacin in diabetic rats. This result showed that the endogenous production of H₂S plays an important protective role against ulceration induced by indomethacin. Non-steroidal anti-inflammatory drugs through inhibiting cyclooxygenases (1 and 2) decrease the production of prostaglandins, increase the rate of gastric acid output and at the same time decrease the rate of mucus secretion (Musumba et al., 2009). As mentioned above, indomethacin by increasing the gastric acid output, induces gastric ulceration. Therefore, multiplying the gastric acid output in diabetic rats due to decreased output of mucosal H₂S in one hand and administration of indomethacin in the other hand, dominated the antacid effect of NaHS at 320 µg/kg.

Conclusion

The results of the present study showed that NaHS effectively decreased the gastric mucosal ulceration induced by indomethacin in diabetic rats through decreasing the gastric acid output and increasing the rate of mucus secretion in the gastric wall of diabetic rats.

Acknowledgments

The source of data used in this paper was from the M.Sc thesis of Miss Nasim Nazariani a student of Payame Noor University, Tehran, Iran. The authors thank to Dr. Eshraf Mansouri for histological evaluation of the results.

Conflict of interest

The authors declare that they have no conflict of interest

References

Boehme MW, Autschbach F, Ell C, Raeth U. Prevalence of silent gastric ulcer, erosions or severe acute gastritis in patients with type 2 diabetes mellitus--a cross-sectional

study. *Hepatogastroenterology* 2007; 54: 643-8.

Goldin E, Ardite E, Elizalde JI, Odriozola A, Panes J, Pique JM, et al. Gastric mucosal damage in experimental diabetes in rats: role of endogenous glutathione. *Gastroenterology* 1997; 112: 855-63.

Kianbakht S, Mozaffari K. Effects of saffron and its active constituents, crocin and safranal, on prevention of indomethacin induced gastric ulcers in diabetic and nondiabetic rats. *J Med Plante* 2009; 8: 30-38.

Lluch I, Ascaso JF, Mora F, Minguez M, Pena A, Hernandez A, et al. Gastroesophageal reflux in diabetes mellitus. *Am J Gastroenterol* 1999; 94: 919-24.

Magierowski M, Jasnos K, Kwiecien S, Drozdowicz D, Surmiak M, Strzalka M, et al. Endogenous prostaglandins and afferent sensory nerves in gastroprotective effect of hydrogen sulfide against stress-induced gastric lesions. *PLoS One* 2015; 10: e0118972.

Mard SA, Ahmadi I, Ahangarpour A, Gharib-Naseri MK, Badavi M. Delayed gastric emptying in diabetic rats caused by decreased expression of cystathionine gamma lyase and h₂ s synthesis: In vitro and in vivo studies. *Neurogastroenterol Motil* 2016; 28: 1677-1689.

Mard SA, Askari H, Neisi N, Veisi A. Antisecretory effect of hydrogen sulfide on gastric acid secretion and the involvement of nitric oxide. *Biomed Res Int* 2014; 2014: 480921.

Mard SA, Mahini S, Dianat M, Farbood Y. Effect of sodium hydrosulfide on mRNA expression of prostaglandin e₂ receptors in response to mucosal acidification and distention-induced gastric acid secretion in rats. *Iran J Basic Med Sci* 2017; 20: 145-149.

Musumba C, Pritchard DM, Pirmohamed M. Review article: cellular and molecular mechanisms of NSAID-induced peptic ulcers. *Aliment Pharmacol Ther* 2009; 30:517-31.

Qiu WC, Wang ZG, Lv R, Wang WG, Han XD, Yan J, et al. Ghrelin improves delayed gastrointestinal transit in alloxan-induced diabetic mice. *World J Gastroenterol* 2008; 14: 2572-7.

Shay H. A simple method for the uniform production of gastric ulceration in rat. *Gastroenterology* 1945; 5:43-61.

Takeuchi K, Ueshima K, Ohuchi T, Okabe S. Induction of gastric lesions and hypoglycemic response by food deprivation in streptozotocin-diabetic rats. *Dig Dis Sci* 1994; 39: 626-34.

Tashima K, Korolkiewicz R, Kubomi M, Takeuchi K. Increased susceptibility of gastric mucosa to ulcerogenic stimulation in diabetic rats--role of capsaicin-sensitive sensory neurons. *Br J Pharmacol* 1998; 124: 1395-402.

Wallace JL. Hydrogen sulfide: a rescue molecule for mucosal defence and repair. *Dig Dis Sci* 2012; 57: 1432-4.